

1	1	2,	1	1	1	3,	3,
	3,	3,	1	4	1	1,5	
(1.			200241	2.		201500;	
3.			200041;	4.		201200	
	5.						

(表 2), 平均监测时长(125.0 ± 105.98) d、平均数据个数($2\,777.3 \pm 2\,509.60$) 个. 9 只监测个体中有 4 只监测到了性行为, 包括 2 只斑嘴鸭 ($2000-2002$) 和 2 只绿头鸭 ($2000-2002$) (表 2). 我们估计最早

Tab. 5 Distribution of GPS locations of each tracked individual in the four types of habitat

- [1] . [J]. 2011 50(3) : 334-343.
- [2] . [J].
2007, 46(6) 906-912.
- [3] .

- [23] HIGUCHI H, OZAKI K, FIJITA G, et al. Satellite tracking of white-naped crane migration and the importance of the Korean demilitarized zone[J]. *Conservation Biology*, 1996,10(3): 806-812.
- [24] . [J]. ,2004(2) : 102.
- [25] , . [J]. 2005(6) : 657-658.
- [26] . (*Grus nigricollis*) [D]. 2006.
- [27] . [J]. 2018, 37(4): 361-372.
- [28] KUANG F, WU W, KE W, et al. Habitat use

Spatial Behavior and Habitat Use of *Anas poecilorhyncha* and *A. platyrhynchos* in Win